

FILE 'REGISTRY' ENTERED AT 16:15:00 ON 09 SEP 2008

L1 STRUCTURE UPLOADED

L2 0 S L1

L3 2 S L1 SSS FULL

FILE 'CAPLUS' ENTERED AT 16:16:24 ON 09 SEP 2008

L4 6 S L3

=> file registry
COST IN U.S. DOLLARS
FULL ESTIMATED COST

SINCE FILE	TOTAL
ENTRY	SESSION
0.21	0.21

FILE 'REGISTRY' ENTERED AT 16:15:00 ON 09 SEP 2008
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2008 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file
provided by InfoChem.

STRUCTURE FILE UPDATES: 8 SEP 2008 HIGHEST RN 1047724-15-1
DICTIONARY FILE UPDATES: 8 SEP 2008 HIGHEST RN 1047724-15-1

New CAS Information Use Policies, enter HELP USAGETERMS for details.


TSCA INFORMATION NOW CURRENT THROUGH July 5, 2008.

Please note that search-term pricing does apply when
conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and
predicted properties as well as tags indicating availability of
experimental property data in the original document. For information
on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stdoc/properties.html>

=>
Uploading C:\Program Files\STNEXP\Queries\10534660alkylmethionine.str



chain nodes :
1 2 3 4 5 6 7 8 9 10 11 12 13
chain bonds :
1-2 2-3 2-10 2-13 3-4 4-5 5-6 5-7 5-9 7-8 10-11 10-12
exact/norm bonds :
2-10 2-13 5-7 5-9
exact bonds :
1-2 2-3 3-4 4-5 5-6 7-8 10-11 10-12

Match level :
1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 6:CLASS 7:CLASS 8:CLASS 9:CLASS
10:CLASS 11:CLASS 12:CLASS 13:CLASS
Generic attributes :
13:
Saturation : Saturated

Element Count :
Node 13: Limited
C,C1-8

L1 STRUCTURE UPLOADED

=> s l1

SAMPLE SEARCH INITIATED 16:15:31 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 12 TO ITERATE

100.0% PROCESSED 12 ITERATIONS

0 ANSWERS

SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
 BATCH **COMPLETE**

PROJECTED ITERATIONS: 33 TO 447


PROJECTED ANSWERS: 0 TO 0

L2 0 SEA SSS SAM L1

=> d l1

L1 HAS NO ANSWERS

L1 STR



Structure attributes must be viewed using STN Express query preparation.

=> s l1 sss full

FULL SEARCH INITIATED 16:16:11 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 169 TO ITERATE

100.0% PROCESSED 169 ITERATIONS

2 ANSWERS


SEARCH TIME: 00.00.01

L3 2 SEA SSS FUL L1

=> d l3 scan

L3 2 ANSWERS REGISTRY COPYRIGHT 2008 ACS on STN


IN Isovaline, 4-(S-methylsulfonylimidoyl)- (9CI)
MF C6 H14 N2 O3 S



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L3 2 ANSWERS REGISTRY COPYRIGHT 2008 ACS on STN
IN Butanoic acid, 2-amino-2-ethyl-4-(S-methylsulfonylimidoyl)- (9CI)
MF C7 H16 N2 O3 S



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

ALL ANSWERS HAVE BEEN SCANNED

=> file caplus
COST IN U.S. DOLLARS
FULL ESTIMATED COST

SINCE FILE	TOTAL
ENTRY	SESSION
178.82	179.03

FILE 'CAPLUS' ENTERED AT 16:16:24 ON 09 SEP 2008
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 9 Sep 2008 VOL 149 ISS 11
FILE LAST UPDATED: 8 Sep 2008 (20080908/ED)

Caplus now includes complete International Patent Classification (IPC) reclassification data for the second quarter of 2008.

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/legal/infopolicy.html>

=> s l3

L4 6 L3

=> d l4 1-6 ti bs bib

'BS' IS NOT A VALID FORMAT FOR FILE 'CAPLUS'

The following are valid formats:

ABS ----- GI and AB
ALL ----- BIB, AB, IND, RE
APPS ----- AI, PRAI
BIB ----- AN, plus Bibliographic Data and PI table (default)
CAN ----- List of CA abstract numbers without answer numbers
CBIB ----- AN, plus Compressed Bibliographic Data
CLASS ----- IPC, NCL, ECLA, FTERM
DALL ----- ALL, delimited (end of each field identified)
DMAX ----- MAX, delimited for post-processing
FAM ----- AN, PI and PRAI in table, plus Patent Family data
FBIB ----- AN, BIB, plus Patent FAM
IND ----- Indexing data
IPC ----- International Patent Classifications
MAX ----- ALL, plus Patent FAM, RE
PATS ----- PI, SO
SAM ----- CC, SX, TI, ST, IT
SCAN ----- CC, SX, TI, ST, IT (random display, no answer numbers;
 SCAN must be entered on the same line as the DISPLAY,
 e.g., D SCAN or DISPLAY SCAN)
STD ----- BIB, CLASS

IABS ----- ABS, indented with text labels
IALL ----- ALL, indented with text labels
IBIB ----- BIB, indented with text labels
IMAX ----- MAX, indented with text labels
ISTD ----- STD, indented with text labels

OBIB ----- AN, plus Bibliographic Data (original)
OIBIB ----- OBIB, indented with text labels

SBIB ----- BIB, no citations
SIBIB ----- IBIB, no citations

HIT ----- Fields containing hit terms
HITIND ----- IC, ICA, ICI, NCL, CC and index field (ST and IT)
 containing hit terms
HITRN ----- HIT RN and its text modification
HITSTR ----- HIT RN, its text modification, its CA index name, and
 its structure diagram
HITSEQ ----- HIT RN, its text modification, its CA index name, its
 structure diagram, plus NTE and SEQ fields
FHITSTR ----- First HIT RN, its text modification, its CA index name, and
 its structure diagram
FHITSEQ ----- First HIT RN, its text modification, its CA index name, its

structure diagram, plus NTE and SEQ fields
 KWIC ----- Hit term plus 20 words on either side
 OCC ----- Number of occurrence of hit term and field in which it occurs

To display a particular field or fields, enter the display field codes. For a list of the display field codes, enter HELP DFIELDS at an arrow prompt (=>). Examples of formats include: TI; TI,AU; BIB,ST; TI,IND; TI,SO. You may specify the format fields in any order and the information will be displayed in the same order as the format specification.

All of the formats (except for SAM, SCAN, HIT, HITIND, HITRN, HITSTR, FHITSTR, HITSEQ, FHITSEQ, KWIC, and OCC) may be used with DISPLAY ACC to view a specified Accession Number.

ENTER DISPLAY FORMAT (BIB):d his
 'D' IS NOT A VALID FORMAT FOR FILE 'CAPLUS'

The following are valid formats:

ABS ----- GI and AB
 ALL ----- BIB, AB, IND, RE
 APPS ----- AI, PRAI
 BIB ----- AN, plus Bibliographic Data and PI table (default)
 CAN ----- List of CA abstract numbers without answer numbers
 CBIB ----- AN, plus Compressed Bibliographic Data
 CLASS ----- IPC, NCL, ECLA, FTERM
 DALL ----- ALL, delimited (end of each field identified)
 DMAX ----- MAX, delimited for post-processing
 FAM ----- AN, PI and PRAI in table, plus Patent Family data
 FBIB ----- AN, BIB, plus Patent FAM
 IND ----- Indexing data
 IPC ----- International Patent Classifications
 MAX ----- ALL, plus Patent FAM, RE
 PATS ----- PI, SO
 SAM ----- CC, SX, TI, ST, IT
 SCAN ----- CC, SX, TI, ST, IT (random display, no answer numbers;
 SCAN must be entered on the same line as the DISPLAY,
 e.g., D SCAN or DISPLAY SCAN)
 STD ----- BIB, CLASS

 IABS ----- ABS, indented with text labels
 IALL ----- ALL, indented with text labels
 IBIB ----- BIB, indented with text labels
 IMAX ----- MAX, indented with text labels
 ISTD ----- STD, indented with text labels

 OBIB ----- AN, plus Bibliographic Data (original)
 OIBIB ----- OBIB, indented with text labels

 SBIB ----- BIB, no citations
 SIBIB ----- IBIB, no citations

 HIT ----- Fields containing hit terms
 HITIND ----- IC, ICA, ICI, NCL, CC and index field (ST and IT)
 containing hit terms
 HITRN ----- HIT RN and its text modification
 HITSTR ----- HIT RN, its text modification, its CA index name, and
 its structure diagram
 HITSEQ ----- HIT RN, its text modification, its CA index name, its
 structure diagram, plus NTE and SEQ fields
 FHITSTR ----- First HIT RN, its text modification, its CA index name, and

its structure diagram
 FHITSEQ ----- First HIT RN, its text modification, its CA index name, its
 structure diagram, plus NTE and SEQ fields
 KWIC ----- Hit term plus 20 words on either side
 OCC ----- Number of occurrence of hit term and field in which it occurs

To display a particular field or fields, enter the display field
 codes. For a list of the display field codes, enter HELP DFIELDS at
 an arrow prompt (=). Examples of formats include: TI; TI,AU; BIB,ST;
 TI,IND; TI,SO. You may specify the format fields in any order and the
 information will be displayed in the same order as the format
 specification.

All of the formats (except for SAM, SCAN, HIT, HITIND, HITRN, HITSTR,
 FHITSTR, HITSEQ, FHITSEQ, KWIC, and OCC) may be used with DISPLAY ACC
 to view a specified Accession Number.
 ENTER DISPLAY FORMAT (BIB):ti abs bib

L4 ANSWER 1 OF 6 CAPLUS COPYRIGHT 2008 ACS on STN
 TI Anti-microbial agents derived from methionine sulfoximine analogues and
 use for treating mycobacterial infections
 AB Novel antimicrobial compns. containing analogs of L-methionine-SR-sulfoximine
 (MSO) that are effective in treating intracellular pathogen infections are
 provided. Specifically, the compns. provided are MSO analogs having
 superior antimicrobial activity with significantly less toxicity as
 compared to MSO. These MSO analogs are suitable for use in treating
 infection in animals including primates, cows, pigs, horses, rabbits,
 mice, rats, cats, and dogs. Moreover, the MSO analogs are ideally suited
 for treating infections caused by the genus Mycobacterium. Addnl.,
 methods for using the novel MSO analogs are also provided.
 AN 2004:452975 CAPLUS <<LOGINID:20080909>>
 DN 141:12262
 TI Anti-microbial agents derived from methionine sulfoximine analogues and
 use for treating mycobacterial infections
 IN Harth, Gunter; Griffith, Owen W.; Horwitz, Marcus A.
 PA Regents of the University of California, USA
 SO PCT Int. Appl., 40 pp.
 CODEN: PIXXD2

DT Patent
 LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004045539	A2	20040603	WO 2003-US36705	20031117
	WO 2004045539	A9	20040805		
	WO 2004045539	A3	20041111		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR				
	AU 2003295579	A1	20040615	AU 2003-295579	20031117
	US 20040157802	A1	20040812	US 2003-715679	20031117
	US 20060142251	A1	20060629	US 2005-534660	20051128
PRAI	US 2002-426502P	P	20021115		
	US 2002-430407P	P	20021202		
	WO 2003-US36705	W	20031117		
OS	MARPAT 141:12262				

L4 ANSWER 2 OF 6 CAPLUS COPYRIGHT 2008 ACS on STN
 TI Pyruvate analog adducts with NAD as lactate dehydrogenase inhibitors
 AB Adducts of pyruvate and NAD⁺ adducts are lactate dehydrogenase inhibitors that can pass through the blood-brain barrier and are of use in the treatment of primary systemic lactic acidosis are prepared and characterized. A series of Na arylidene pyruvates were prepared and the adducts with NAD⁺ prepared by standard chemical. These were then tested for inhibition of beef heart and rat brain lactate dehydrogenases. An NAD-pyruvate reduced the activity of the beef heart enzyme to 90% of control values and reduced the activity of the rat brain enzyme to 48% of controls in the presence of 0.24 mM pyruvate. An aldehyde analog was similarly active in the nanomolar range. Inhibition of lactate dehydrogenase activity in synaptosomes was also demonstrated.

AN 1991:38443 CAPLUS <<LOGINID::20080909>>

DN 114:38443

OREF 114:6623a,6626a

TI Pyruvate analog adducts with NAD as lactate dehydrogenase inhibitors

IN Cooper, Arthur J. L.

PA Cornell Research Foundation, Inc., USA

SO U.S., 8 pp.

CODEN: USXXAM

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 4950602	A	19900821	US 1987-16894	19870220
PRAI	US 1987-16894		19870220		
OS	MARPAT 114:38443				

L4 ANSWER 3 OF 6 CAPLUS COPYRIGHT 2008 ACS on STN

TI Amino acid sulfoximines: α -ethylmethionine sulfoximine

AB α -Ethylmethionine sulfoximine, HO₂CCet(NH₂)CH₂CH₂S(O)Me:NH, was prepared by treatment of HO₂CCet(NH₂)CH₂CH₂SMe (I) with HCl. I was prepared by treatment of EtCOCH:CH₂ with MeSH to give EtCOCH₂CH₂SMe which was converted to a hydantoin derivative with (NH₄)₂CO₃ and NaCN and the product hydrolyzed to I.

AN 1988:132274 CAPLUS <<LOGINID::20080909>>

DN 108:132274

OREF 108:21719a,21722a

TI Amino acid sulfoximines: α -ethylmethionine sulfoximine

AU Griffith, Owen W.

CS Med. Coll., Cornell Univ., New York, NY, 10021, USA

SO Methods in Enzymology (1987), 143(Sulfur Sulfur Amino Acids), 286-91

CODEN: MENZAU; ISSN: 0076-6879

DT Journal

LA English

L4 ANSWER 4 OF 6 CAPLUS COPYRIGHT 2008 ACS on STN

TI Inhibition of glutathione biosynthesis by prothionine sulfoximine

(S-n-propyl homocysteine sulfoximine), a selective inhibitor of

γ -glutamylcysteine synthetase

AB DL-Prothionine SR-sulfoximine [70085-86-8] and α -methyl-DL-prothionine-SR-sulfoximine [70056-05-2] were prepared and found to markedly inhibit γ -glutamylcysteine synthetase [9023-64-7] but to not significantly affect glutamine synthetase [9023-70-5]. After injection of prothionine sulfoximine into mice, the level of kidney glutathione [70-18-8] decreased rapidly to .apprx.20% of the control level indicating that a large fraction, rather than a small pool, of glutathione participates in rapid turnover. The rapid decline of the glutathione

level that occurs after inhibition of glutathione synthesis reflects the normal rate of intracellular glutathione utilization by the γ -glutamyl cycle. A number of related sulfoximines were synthesized and tested as inhibitors of glutamine and γ -glutamylcysteine synthetases.

AN 1979:198299 CAPLUS <<LOGINID::20080909>>

DN 90:198299

OREF 90:31455a,31458a

TI Inhibition of glutathione biosynthesis by prothionine sulfoximine (S-n-propyl homocysteine sulfoximine), a selective inhibitor of γ -glutamylcysteine synthetase

AU Griffith, Owen W.; Anderson, Mary E.; Meister, Alton

CS Med. Coll., Cornell Univ., New York, NY, USA

SO Journal of Biological Chemistry (1979), 254(4), 1205-10

CODEN: JBCHA3; ISSN: 0021-9258

DT Journal

LA English

L4 ANSWER 5 OF 6 CAPLUS COPYRIGHT 2008 ACS on STN

TI Differential inhibition of glutamine and γ -glutamylcysteine synthetases by α -alkyl analogs of methionine sulfoximine that induce convulsions

AB α -Methyl-DL-methionine (SR)-sulfoximine [66735-67-9] and α -ethyl-DL-methionine (SR)-sulfoximine [66735-68-0], like L-methionine (SR)-sulfoximine [15985-39-4], induced convulsions in mice and inhibited glutamine synthetase [9023-70-5] irreversibly; α -ethylmethionine sulfoximine was approx.50% as inhibitory as methionine sulfoximine and α -methylmethionine sulfoximine. However, whereas α -methylmethionine sulfoximine and methionine sulfoximine inhibited γ -glutamylcysteine synthetase [9023-64-7] markedly, α -ethylmethionine sulfoximine did not, nor did administration of the α -Et analog produce the decrease in tissue glutathione [70-18-8] levels found after giving methionine sulfoximine or its α -Me analog. The α -alkyl methionine sulfoximine analogs cannot be catabolized via the corresponding α -keto or α -imino acids, and, like other α -substituted amino acids, are probably not metabolized to a significant extent in vivo; this suggests that the amino acid sulfoximine mols. themselves, rather than their metabolites, are directly involved in the induction of convulsions. Possible explanations for the reported lack of correlation between the occurrence of convulsions and the levels of glutamine synthetase activity (and its substrates and product) are considered.

AN 1978:500916 CAPLUS <<LOGINID::20080909>>

DN 89:100916

OREF 89:15375a,15378a

TI Differential inhibition of glutamine and γ -glutamylcysteine synthetases by α -alkyl analogs of methionine sulfoximine that induce convulsions

AU Griffith, Owen W.; Meister, Alton

CS Dep. Biochem., Cornell Univ. Med. Coll., New York, NY, USA

SO Journal of Biological Chemistry (1978), 253(7), 2333-8

CODEN: JBCHA3; ISSN: 0021-9258

DT Journal

LA English

L4 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2008 ACS on STN

TI Sulfur-containing amino acids

GI For diagram(s), see printed CA Issue.

AB MeCH:CHCHO (140 g.) and 96 g. MeSH in the presence of 2 drops of piperidine stirred 0.5 hr. at 5-10° and 3 hrs. at room temperature, the mixture treated with an addnl. 28 g. MeSH, heated about 1 hr. at 90°,

diluted with 500 cc. Et₂O, washed with dilute HCl and H₂O, dried, and evaporated, and the residue distilled gave 201 g. MeSCHMeCH₂CHO (I), b₂₃ 80°. AcCH:CH₂ (27 g.) and 18 g. MeSH yielded 35.4 g. Ac(CH₂)₂SMe, b₅₅ 106°, n_{D25} 1.4711. I (48.5 g.), 113 g. (NH₄)₃SO₃, 25.5 g. NaCN, 335 cc. EtOH, and 335 cc. H₂O heated 5 hrs. with stirring at 55°, the mixture concentrated to about 300 cc., treated cautiously with 50 cc. concentrated

HCl, heated 7 min. at about 90°, refrigerated, and filtered, and the residue washed with 200 cc. H₂O yielded 49 g. 5-(β-benzylmercapto)propylhydantoin, m. 117-18° (from EtOAc). Similarly were prepared the following compds. RR'C.CO.NH.CO.NH (R, R', m.p., and % yield given): MeS(CH₂)₂, Me, 109.5-10.5°, 93.8; MeSCHMeCH₂, H, 191-2°, 50.1; MeSCHPhCH₂, H, 173-4°, 491. S-Benzyl-4-methylhomocysteine (7.17 g.), m. 222.5-3.5° (decomposition) (from H₂O) (obtained in 94% yield from the hydantoin) (0.69, 0.74, 0.93) (the figures given in parentheses through out this abstract represent the R_f values of the resp. compds. obtained by ascending paper chromatography with BuOH-AcOH, lutidine-collidine, and PhOH-H₂O, resp.) in 300 cc. liquid NH₃ treated with about 1.7 g. Na, the solution decolorized with about 1 g. NH₄Cl, treated with 5 cc. MeI, and evaporated, the residue treated with 125 cc. H₂O, washed with Et₂O, filtered, neutralized with concentrated HCl to pH about 6, concentrated to about 50 cc., diluted with 50 cc. Me₂CO, and refrigerated, and the crystalline deposit recrystd. from aqueous MeOH yielded

4.1 g. MeSCHMeCH₂CH(NH₂)CO₂H (II), m. 236-7° (decomposition), (0.44, 0.53, 0.79). Similarly were prepared: MeS(CH₂)₂CMe(NH₂)CO₂H, 61%, m. 284-5° (decomposition) (from aqueous MeOH), (0.45, 0.50, 0.77); MeSCHPh(CH₂)₂CH(NH₂)CO₂H, 49.3%, m. 201-2° (decomposition) (from H₂O). BzCH₂SMe (21.8 g.) in 50 cc. dry Et₂O added with stirring to 1.4 g. LiAlH₄ in 10 cc. dry Et₂O, the mixture refluxed 1 hr. with stirring, cooled, and treated with stirring with 200 cc. ice water and 100 cc. 5N H₂SO₄, the aqueous layer washed with Et₂O, the combined Et₂O solns. washed, dried, and evaporated under a jet of dry air, and the residue distilled gave 18.4 g. MeSCH₂CH(OH)Ph (III), b_{1.8} 113-14.5°. III (170 mg.) treated with MeI yielded III. MeI, m. 134-5° (decomposition). III (15.8 g.) in 25 cc. dry CHCl₃ treated with cooling with 9.2 g. SOCl₂ in 15 cc. dry CHCl₃, the mixture cooled 0.5 hr., kept at room temperature overnight and evaporated, the residue heated gently with 5 cc. dry CHCl₃ and 5 cc. SOCl₂, and the mixture distilled gave 14.3 g. MeSCH₂CHClPh (IV), b_{2.8} 106-7°, n_{D25} 1.5692. AcNHCH(CO₂Et)₂ (11.6 g.) and 200 mg. KI added with stirring to 1.23 g. Na in 100 cc. absolute EtOH, the mixture treated with 10 g. IV in 1 portion, stirred 2 hrs. at room temperature, refluxed 5 hrs., and filtered hot, the residue washed with about 50 cc. hot EtOH, the combined alc. solns. evaporated to dryness in vacuo, the residual oil kept at room temperature overnight, and the crystalline material washed with dilute HCl and H₂O and dried in vacuo over KOH pellets yielded 16 g. MeSCH₂CHPhC(NHAc)(CO₂Et)₂ (V), m. 95-6° (from Et₂O-pentane). Crude V (14.4 g.), 40 cc. H₂O, and 10 cc. concentrated

HCl refluxed 6 hrs. with stirring, the mixture treated with 40 cc. H₂O and 10 cc. concentrated HCl, refluxed 1.5 hrs. with stirring, cooled to room temperature, the solid refluxed 8 hrs. with stirring with 80 cc. glacial AcOH and 10 cc. concentrated HCl, treated with Norit, and filtered, the residue washed with

H₂O, the combined filtrates evaporated in vacuo, the residue (about 10 g.) triturated with 50 cc. Me₂CO and filtered, and the residue washed with Me₂CO and dried yielded 5 g. MeSCH₂CHPhC(NH₂)CO₂H·HCl (VI·HCl), m. 208-9° (decomposition); the Me₂CO solns. combined and evaporated to dryness, the residue refluxed 6.5 hrs. with 25 cc. H₂O, 25 cc. glacial AcOH, and 10 cc. concentrated HCl, the solution evaporated to dryness in vacuo, the residue washed

with Me₂CO and neutralized with AmNH₂, and a 1-g. portion dissolved in 8 cc. H₂O and neutralized with AmNH₂ to pH 6, diluted with 25 cc. Me₂CO, and filtered, and the residue washed with 15 cc. Me₂CO yielded 300 mg. VI; the filtrate diluted with Me₂CO gave a 2nd crop, 350 mg. MeSH (14 g.) passed with stirring and cooling into 1.2 g. Na in 150 cc. absolute MeOH, the mixture treated with stirring and cooling with 50 g. Me α -benzamidosenecioate, diluted with 200 cc. absolute MeOH and 200 cc. dry C₆H₆, stirred 1 hr. at room temperature, allowed to stand overnight, treated with

3.12

g. glacial AcOH, and evaporated to dryness in vacuo at room temperature, the residue

washed with warm dry C₆H₆, the C₆H₆ evaporated, the residue (58 g.), 300 cc. 85% HCO₂H, 300 cc. concentrated HCl, and 300 cc. H₂O refluxed 6 hrs., the

solution

concentrated to about 50 cc., washed with Et₂O, neutralized with AmNH₂ to pH 6, diluted with 350 cc. Me₂CO, and refrigerated 2 days, and the white crystals washed with 300 cc. Me₂CO and 200 cc. Et₂O yielded 16.8 g.

S-methylpenicillamine, m. 281-2° (0.38, 0.50, 0.80); it was also obtained in the same manner from 2-phenyl-4-isopropylidene-5-oxazolone and 30 g. MeSH. MeSH (16 g.) passed into 1.2 g. Na in 300 cc. absolute MeOH, the solution treated with cooling and stirring with 62.3 g. 2-phenyl-4-benzal-5-oxazolone in 500 cc. warm, dry C₆H₆, the mixture stirred about 1 hr., kept at room temperature, treated with 3.12 g. glacial AcOH, and evaporated to

dryness in

vacuo, the residue treated with 100 cc. warm C₆H₆ and filtered, the filtrate diluted with 100 cc. warm C₆H₆ and 500 cc. pentane, and chilled, and the deposit washed with 150 cc. pentane yielded 74 g.

PhCH(SMe)CH(NHBz)CO₂Me (VII), m. 97-8.5° (from EtOAc-pentane).

Crude VII (32.9 g.) hydrolyzed with 150 cc. H₂O, 150 cc. concentrated HCl, and 150 cc. 90% HCO₂H, the solution concentrated in vacuo to near dryness, and the

precipitate

washed with three 100-cc. portions H₂O, dissolved in 75 cc. H₂O, neutralized to pH 6 with AmNH₂, and chilled yielded 12.5 g. S-methyl-3-phenylcysteine, m. 178-9° (decomposition) (0.51, 0.65, 0.88).

The following sulfoxides were prepared by oxidation of the appropriate sulfides with H₂O₂ by the method of Toennies and Kolb (C.A. 33, 5359.9) (% yield, m.p., and Rf values given): PhCH₂S(O)CHMeCH₂CH(NH₂)CO₂H, 64.7, 214-15° (decomposition) (from H₂O), (0.45, 0.60, 0.92);

MeS(O)CH₂CH₂CM₂(NH₂)CO₂H, 91.8, 239.5-40.5° (decomposition) (from aqueous MeOH), (0.14, 0.35, 0.77); MeS(O)CHMeCH₂CH(NH₂)CO₂H (VIII), 84.4, 213.5-14.5° (from aqueous MeOH), (0.13, 0.40, 0.80);

MeS(O)CH₂CHPhCH(NH₂)CO₂H, 74.4, 205-6° (decomposition) (from aqueous MeOH), (0.33, 0.59, 0.87); MeS(O)CHPhCH₂CH(NH₂)CO₂H, 87.7, 189-90° (decomposition) (from aqueous MeOH), (0.33, 0.47, 0.85);

Me₂CHCH[S(O)Me]CH(NH₂)CO₂H, 77.7, 166-7° (from aqueous MeOH), (0.14, 0.40, 0.76); PhCH[S(O)Me]CH(NH₂)CO₂H, 73.2, 147-8° (decomposition) (from aqueous MeOH), (0.29, 0.54, 0.82). VIII (600 mg.), 3 cc. H₂O, 2 cc. MeOH, 0.2 cc. concentrated HCl, and 2 cc. 30% H₂O₂ refluxed 2 hrs., treated with 1 cc.

30%

H₂O₂, refluxed again 2 hrs., neutralized with AmNH₂ to pH 6.5, diluted with 100 cc. Me₂CO and filtered, and the residue washed with 50 cc. Me₂CO yielded 550 mg. MeS(O₂)CHMeCH₂CH(NH₂)CO₂H, m. 230-1° (decomposition) (from aqueous MeOH), (0.14, 0.50, 0.72). In the same manner was prepared PhCH₂S(O₂)CH₂CH₂CH(NH₂)CO₂H, 70.6%, m. 229-30° (decomposition) (from H₂O), (0.50, 0.65, 0.84). The following sulfones were prepared by the oxidation on the appropriate sulfides with H₂O₂ in the presence of NH₄ molybdate and HClO₄ by the method of Toennies and Kolb (C.A. 35, 6571.1) (% yield, m.p., and Rf values given): MeS(O₂)CH₂CH₂CM₂(NH₂)CO₂H, 73.6, 288-9° (decomposition) (from aqueous MeOH), (0.16, 0.45, 0.65); MeS(O₂)CH₂CHPhCH(NH₂)CO₂H (IX), 50.8, 222-3° (decomposition) (from H₂O), (0.32, 0.61, 0.79); MeS(O₂)CHPhCH₂CH(NH₂)CO₂H (X), 95.4, 196.5-7.5°

(decomposition), (0.37, 0.55, 0.79); Me₂CHCH[S(O₂)Me]CH(NH₂)CO₂H, 77.7, 166-7° (from aqueous MeOH), (0.14, 0.53, 0.68); MeS(O₂)CHPhCH(NH₂)CO₂H, 51.2, 141-2° (decomposition) (from aqueous MeOH), (0.30, 0.52, 0.70). VIII (6.0 g.) treated dropwise with stirring at 3° with 10.4 cc. concentrated H₂SO₄, the mixture heated with stirring to 45°, treated during 1 hr. at 48° with 54 cc. 1.4N HN₃ in CHCl₃, then heated with stirring 5 hrs. at 48°, treated with 13.5 cc. HN₃ solution, heated 5 hrs. with stirring at 50°, stirred overnight at room temperature, poured with stirring onto 75 g. crushed ice, neutralized with solid Ba(OH)₂ to about pH 2.5 then to pH 5 with solid BaCO₃, and centrifuged, the supernatant decanted, the residue mixed with H₂O, centrifuged, and decanted, this operation repeated until free of amino acid, the combined aqueous solns. concentrated in vacuo at 50° to about 100 cc., treated with C, and filtered, and the filtrate concentrated to about 40 cc., filtered, and evaporated to dryness yielded 6.4 g. MeS(:NH)CHMeCH₂CH(NH₂)CO₂H, m. 199-200° (decomposition) (from aqueous MeOH), (0.08, 0.38, 0.71). In the same manner was prepared: MeS(:NH)CH₂CH₂CHMe(NH₂)CO₂H, 100, 199-200° (decomposition) (from aqueous MeOH), (0.10, 0.35, 0.67). IX (100 mg.) treated with about 60 mg. N-bromosuccinimide gave MeS(O₂)CH₂CHPhCHO, isolated as the 2,4-dinitrophenylhydrazone, m. 188-9° (decomposition). X gave similarly MeS(O₂)CHPhCH₂CHO, isolated as the 2,4-dinitrophenylhydrazone, decomposed at 196-8° with a change from yellow to red at 169°. Only 4 of the amino acids suppressed the multiplication of T2 bacteriophage of Escherichia coli strain A.T.C.C. number 11303 at pH 7 and 37° at 100 p.p.m. or less.

AN 1956:73727 CAPLUS <<LOGINID::20080909>>
DN 50:73727
OREF 50:13802g-i,13803a-i,13804a-g
TI Sulfur-containing amino acids
AU Reisner, David B.
CS Wallace & Tiernan, Inc., Newark, NJ
SO Journal of the American Chemical Society (1956), 78, 2132-5
CODEN: JACSAT; ISSN: 0002-7863
DT Journal
LA Unavailable
OS CASREACT 50:73727